


64
CLAIMS

1. Benzopyran derivatives of the general formula



(I)

wherein :

D represents S or O;

- 10 R_1 , R_2 , R_3 and R_4 are independently hydrogen, halogen, C_{1-6} -alkyl, C_{3-8} -cycloalkyl, hydroxy, C_{1-6} -alkoxy, C_{1-6} -alkoxy- C_{1-6} -alkyl, nitro, amino, cyano, cyanomethyl, perhalomethyl, C_{1-6} -monoalkyl- or dialkylamino, sulfamoyl, C_{1-6} -alkylthio, C_{1-6} -alkylsulfonyl, C_{1-6} -alkylsulfinyl, formyl, C_{1-6} -alkylcarbonylamino, R_8 arylthio, R_8 arylsulfinyl, R_8 arylsulfonyl, C_{1-6} -alkoxycarbonyl, C_{1-6} -alkoxycarbonyl- C_{1-6} -alkyl, carbamoyl, carbamoylmethyl, C_{1-6} -monoalkyl- or dialkylaminocarbonyl, C_{1-6} -monoalkyl- or dialkylaminothiocarbonyl, ureido, C_{1-6} -monoalkyl- or dialkylaminocarbonylamino, thioureido, C_{1-6} -monoalkyl- or dialkylaminothiocarbonylamino, C_{1-6} -monoalkyl- or dialkylaminosulfonyl, carboxy, carboxy- C_{1-6} -alkyl, acyl, R_8 aryl, R_8 aryl- C_{1-6} -alkyl, R_8 aryloxy;

20

R_5 and R_6 are each independently hydrogen, C_{1-6} -alkyl or, R_5 and R_6 taken together with the carbon atom to which they are attached form a 3- to 6-membered carbocyclic ring;

25

65

R₇ is 2-, 3- or 4-pyridyl optionally mono- or polysubstituted by R₁ or

R₇ is 2- or 3-thienyl optionally mono- or polysubstituted substituted by R₁ or

R₇ is phenyl mono- or polysubstituted by R₁ with the exception of R₇ representing C₆H₅;

5

R₈ is hydrogen, halogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, hydroxy, C₁₋₆-alkoxy, nitro, amino, cyano, cyanomethyl, perhalomethyl;

10 or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture or any polymorphic and tautomeric form.

2. A benzopyran derivative according to claim 1 wherein D represents S.

15

3. A benzopyran derivative according to claim 1 or 2 selected from :

R/S-4-(3-Chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,

20 R/S-6-Chloro-4-(3-chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,

R/S-4-(4-Chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,

25 R/S-6-Chloro-4-(4-chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,

R/S-6-Bromo-4-(4-chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,

R/S-4-(3-Cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,

30 R/S-6-Chloro-4-(3-cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,

- R/S-6-Bromo-4-(3-cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-4-(4-Cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,
5 R/S-6-Chloro-4-(4-cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-6-Bromo-4-(4-cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(3-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
10 R/S-6-Bromo-3,4-dihydro-2,2-dimethyl-4-(3-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
R/S-3,4-Dihydro-2,2-dimethyl-6-fluoro-4-(4-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
15 R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(4-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
R/S-4-(3-Chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,
R/S-6-Chloro-4-(3-chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
20 R/S-4-(4-Chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,
R/S-6-Chloro-4-(4-chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
25 R/S-6-Bromo-4-(4-chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-4-(3-Cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,
R/S-6-Chloro-4-(3-cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
30

- R/S-6-Bromo-4-(3-cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-4-(4-Cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,
5 R/S-6-Chloro-4-(4-cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-6-Bromo-4-(4-cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(3-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
10 R/S-6-Bromo-3,4-dihydro-2,2-dimethyl-4-(3-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
R/S-3,4-Dihydro-2,2-dimethyl-6-fluoro-4-(4-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
15 R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(4-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
R/S-4-(3-Chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,
R/S-6-Chloro-4-(3-chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
20 R/S-4-(4-Chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,
R/S-6-Chloro-4-(4-chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
25 R/S-6-Bromo-4-(4-chlorophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-4-(3-Cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,
R/S-6-Chloro-4-(3-cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
30

- R/S-6-Bromo-4-(3-cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-4-(4-Cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,
5 R/S-6-Chloro-4-(4-cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-6-Bromo-4-(4-cyanophenylaminothiocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(3-
10 nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
R/S-6-Bromo-3,4-dihydro-2,2-dimethyl-4-(3-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
R/S-3,4-Dihydro-2,2-dimethyl-6-fluoro-4-(4-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
15 R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(4-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
R/S-6-Bromo-3,4-dihydro-2,2-dimethyl-4-(4-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
R/S-6-Bromo-3,4-dihydro-2,2-dimethyl-4-(4-nitrophenylaminothiocarbonylamino)-2*H*-1-benzopyran,
20 R/S-6-Bromo-3,4-dihydro-2,2-dimethyl-4-(3-trifluoromethylphenylaminothiocarbonylamino)-2*H*-1-benzopyran,
R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(2-methoxyphenylaminocarbonylamino)-2*H*-1-benzopyran,
25 R/S-6-Bromo-3,4-dihydro-2,2-dimethyl-4-(2-methoxyphenylaminocarbonylamino)-2*H*-1-benzopyran,
R/S-3,4-Dihydro-2,2-dimethyl-6-fluoro-4-(3-methoxyphenylaminocarbonylamino)-2*H*-1-benzopyran,
R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(3-methoxyphenylaminocarbonylamino)-2*H*-1-benzopyran,
30 R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(3-methoxyphenylaminocarbonylamino)-2*H*-1-benzopyran,

68

- R/S-6-Bromo-3,4-dihydro-2,2-dimethyl-4-(3-methoxyphenylaminocarbonylamino)-2*H*-1-benzopyran,
R/S-3,4-Dihydro-2,2-dimethyl-6-fluoro-4-(4-methoxyphenylaminocarbonylamino)-2*H*-1-benzopyran,
5 R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(4-methoxyphenylaminocarbonylamino)-2*H*-1-benzopyran,
R/S-6-Bromo-3,4-dihydro-2,2-dimethyl-4-(4-methoxyphenylaminocarbonylamino)-2*H*-1-benzopyran,
R/S-3,4-Dihydro-2,2-dimethyl-6-fluoro-4-(2-methylphenylaminocarbonylamino)-2*H*-1-benzopyran,
10 R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(2-methylphenylaminocarbonylamino)-2*H*-1-benzopyran,
R/S-6-Bromo-3,4-dihydro-2,2-dimethyl-4-(2-methylphenylaminocarbonylamino)-2*H*-1-benzopyran,
15 R/S-3,4-Dihydro-2,2-dimethyl-6-fluoro-4-(3-methylphenylaminocarbonylamino)-2*H*-1-benzopyran,
R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(3-methylphenylaminocarbonylamino)-2*H*-1-benzopyran,
R/S-6-Bromo-3,4-dihydro-2,2-dimethyl-4-(3-methylphenylaminocarbonylamino)-2*H*-1-benzopyran,
20 R/S-3,4-Dihydro-2,2-dimethyl-6-fluoro-4-(4-methylphenylaminocarbonylamino)-2*H*-1-benzopyran,
R/S-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(4-methylphenylaminocarbonylamino)-2*H*-1-benzopyran,
25 R/S-6-Bromo-3,4-dihydro-2,2-dimethyl-4-(4-methylphenylaminocarbonylamino)-2*H*-1-benzopyran,
R/S-4-(2-Chlorophenylaminocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,
R/S-6-Chloro-4-(2-chlorophenylaminocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
30

- R/S-6-Bromo-4-(2-chlorophenylaminocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-4-(3-Chlorophenylaminocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-fluoro-2*H*-1-benzopyran,
5 R/S-6-Chloro-4-(3-chlorophenylaminocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-6-Bromo-4-(3-chlorophenylaminocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-4-(4-Chlorophenylaminocarbonylamino)-3,4-dihydro-2,2-dimethyl-6-
10 fluoro-2*H*-1-benzopyran,
R/S-6-Chloro-4-(4-chlorophenylaminocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran,
R/S-6-Bromo-4-(4-chlorophenylaminocarbonylamino)-3,4-dihydro-2,2-dimethyl-2*H*-1-benzopyran.
15
4. Benzopyran derivatives according to any one of the preceding claims for use as openers of the K_{ATP}-regulated potassium channels.
5. A pharmaceutical composition comprising a benzopyran derivative according
20 to any one of the preceding claims or pharmaceutically acceptable salt thereof with a pharmaceutically acceptable acid or base or any optical isomer or mixture of optical isomers, including a racemic mixture or any tautomeric form together with one or more pharmaceutically acceptable carriers or diluents.
- 25 6. A pharmaceutical composition for use in the treatment of diseases of the endocrinological system such as hyperinsulinaemia and diabetes comprising a benzopyran derivative according to any one of the preceding benzopyran derivative claims or a pharmaceutical acceptable salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of

optical isomers, including a racemic mixture, or any tautomeric form together with a pharmaceutically acceptable carrier or diluent.

5 7. The pharmaceutical composition according to any one of the claims 5 or 6 in the form of an oral dosage unit or parental dosage unit.

10 8. A pharmaceutical composition according to any one of the claims 5 or 6 wherein said benzopyran derivative is administered as a dose in a range from about 0.05 to 1000, preferably from about 0.1 to 500 and especially in the range from 50 to 200 mg per day.

15 9. A benzopyran derivative according to any one of the preceding benzopyran derivative claims or a pharmaceutically acceptable salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric form for therapeutic use.

20 10. A benzopyran derivative according to any one of the preceding benzopyran derivative claims or a pharmaceutically acceptable salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric form for therapeutic use in the treatment of diseases of the endocrinological system, such as hyperinsulinaemia and diabetes.

25 11. The use of a benzopyran derivative according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric form as a medicament.

30

71

12. The use of a benzopyran derivative according to any of the preceding compound claims for preparing a medicament.


13. The use of a benzopyran derivative according to any one of the preceding benzopyran derivative claims or a pharmaceutically acceptable salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric form for the preparation of a medicament for the treatment of diseases of the endocrinological system, such as hyperinsulinaemia and diabetes.

14. A method of treating diseases of the endocrinological system, such as hyperinsulinaemia and diabetes in a subject in need thereof comprising administering an effective amount of a benzopyran derivative according to any one of the preceding benzopyran derivative claims to said subject.

15. A process for the manufacture of a medicament, particular to be use in the treatment of diseases of the endocrinological system, such as hyperinsulinaemia and diabetes which process comprising bringing a compound of formula (I) according to any one of the preceding compound claims 1 or a pharmaceutically acceptable salt thereof into a galenic dosage form.

16. A method of preparing a benzopyran derivative of formula (I) which comprises :

- reacting a compound of formula (II)



(II)

72

wherein R represents NH_2 and R_1 , R_2 , R_3 and R_4 are defined as for formula (I) with an isothiocyanate of formula (III)



wherein D represents S or O and R_7 is defined as for formula (I), to form a benzopyran derivative of formula (I); or

10

- reacting a compound of formula (II) wherein R represents $\text{-N}=\text{C}=\text{S}$ and R_1 , R_2 , R_3 and R_4 are defined as for formula (I) with an amine of formula (IV)



wherein R_7 is defined as for formula (I), to form a benzopyran derivative of formula (I).